



FUGITIVE EMISSIONS

DEVELOPMENT: IPPC PERMIT APPLICATION FOR THE HOUSING OF 260,000 BROILERS

LOCATION: Dunnimere Farm
Portway Lane
Harlaston
Near Tamworth
Staffordshire
B79 9LA

CLIENT: Dunnimere Poultry Ltd

Roger Parry & Partners LLP Carmarthen
C/O The Estates Office, 20 Salop Road, Oswestry, Shropshire, SY11 2NU
Tel: 01691655334 Fax: 01691 657798
Email: gail@rogerparry.net

Also at: 11 Severn Street, Welshpool, Powys, SY21 7AG
Tel: 01938 554499 Fax: 01938 554462
email: welshpool@rogerparry.net

Also at: Hogstow Hall, Minsterley, Shrewsbury, SY5 0HZ
Tel: 01743 791336 Fax: 01743 792770
email: mail@rogerparry.net

Hazard	Receptor	Pathway	Risk Management	Exposure	Consequence	Overall Risk
To Air						
Dust: Sources: Litter and Feed.	Neighbouring dwelling houses within 400m of installation Surrounding Land and Vegetation	Air	Use of suitable bedding materials. Use of pelleted feed delivered in sealed systems. Litter removed carefully during cleanout minimising dust. Full trailers sheeted before leaving.	Dust could have the potential to reach nearby neighbours and surrounding land during certain weather conditions and operations (clean out approximately 14 days per year) Careful management Should prevent this happening. Unlikely during laying phase.	Nuisance – dust on surrounding vegetation, cars and clothing. Smothering and direct damage to nearby vegetation.	Not significant if carefully managed.
Ammonia: Source: Poultry housing and Litter storage	Neighbouring dwelling houses within 400m of installation Surrounding Land and Vegetation	Air	Measures as described in SGN EPR6.09 How to Comply. Litter kept dry and friable. Feed formulated to match flock requirements. Litter removed off site following crop depletion, no storage on site.	The impact of Ammonia Air emissions have been assessed using the H1 methodology and modelling demonstrates there will be some impact to nearby wildlife sites.	Arial deposition and direct toxic effect on trees. Nutrient enrichment of soils and changes to sensitive ecosystems.	Significant reduction demonstrated.
Zoonoses and Notifiable diseases	Human Health an Livestock Health	Air, Direct contact	Detailed biosecurity measures in place. Visitor's procedure. Use of appropriate PPE Tailored terminal hygiene programme Veterinarian health plan	Unlikely	Human and Livestock health implications	Not significant if carefully managed.
To Water						
Wash water run off to nearby ditch	Drainage ditches	Land	Wash water from poultry houses directed in sealed system to underground storage tank. Spillages of litter on yard areas during cleanout swept up, Lightly contaminated yard wash directed to underground tank. All	Unlikely	Pollution of watercourses leading to eutrophication and poisoning of flora and fauna	Not significant if managed carefully.

			clean site drainage directed to soakaways.			
Pests						
Flies	Neighbouring dwelling houses within 400m of installation	Air	Temporary field heaps regularly checked for maggots and flies, heaps treated with pesticide and covered if flies become a an issue	Unlikely	Flies are a vector of pollution that can harm human health and amenity causing offence.	Not significant if managed carefully.
Rodents/Vermin	Neighbouring dwelling houses within 400m of installation	Land	Feed spillages cleared up promptly. Specialist contractor used to control pests.	Unlikely	Rodents are a vector of pollution that can harm human health and amenity causing offence.	Not significant if managed carefully.