



FUGUTIVE EMISSIONS

DEVELOPMENT: IPPC PERMIT VARIATION - EPR/KP3130HL/V005

LOCATION: Great Ness Poultry Farm
Rodefern Lane
Nesscliffe
Shrewsbury
Shropshire
SY4 1AT

CLIENT: Great Ness Poultry Limited

Roger Parry & Partners LLP
Mercian House, 9 Darwin Court, Oxon Business Park, Shrewsbury, SY3 5AL
Tel: 01691655334
Email: richard@rogerparry.net emily@rogerparry.net

Reviewed: December 2025

Fugitive Emissions Assessment & Management Plan

There are four properties within 400 metres of Great Ness Poultry (not including receptors owned by operator/operator staff):

NAME	EASTING	NORTHING	Distance from the additional units (m)	Orientation of receptor from installation
Rodefern Farm, Rodefern Lane, Montford Bridge, SHREWSBURY, SY4 1AT	340240	318357	115m	South-West
Ness Hall, Great Ness, NESSCLIFFE, Shropshire, SY4 2LB	339975	318894	331m	North
Oakfield, Great Ness, SHREWSBURY, Shropshire, SY4 2LB	340048	318890	292m	North

Hazard To Air	Receptor	Pathway	Risk Management	Exposure	Consequence	Overall Risk
Dust: Sources: Litter and Feed.	Neighbouring dwelling houses within 400m of installation Surrounding Land and Vegetation	Air	Use of suitable bedding materials. Use of pelleted feed delivered in sealed systems. Litter removed carefully during cleanout minimising dust. Full trailers sheeted before leaving.	Dust could have the potential to reach nearby neighbours and surrounding land during certain weather conditions and operations (clean out approximately 14 days per year) Careful management Should prevent this happening. Unlikely during laying phase.	Nuisance – dust on surrounding vegetation, cars and clothing. Smothering and direct damage to nearby vegetation.	Not significant if carefully managed.
Ammonia: Source: Poultry housing and Litter storage	Neighbouring dwelling houses within 400m of installation	Air	Measures as described in SGN EPR6.09 How to Comply. Litter kept dry and friable. Feed	The impact of Ammonia Air emissions have been assessed using the H1 methodology and modelling	Arial deposition and direct toxic effect on trees. Nutrient enrichment of soils and	Screened out.

	Surrounding Land and Vegetation		formulated to match flock requirements. Litter removed off site following crop depletion, no storage on site.	demonstrates there will be not be a significant impact to nearby wildlife sites.	changes to sensitive ecosystems.	
Zoonoses and Notifiable diseases	Human Health an Livestock Health	Air, Direct contact	Detailed biosecurity measures in place. Visitor's procedure. Use of appropriate PPE Tailored terminal hygiene programme Veterinarian health plan	Unlikely	Human and Livestock health implications	Not significant if carefully managed.
To Water						
Wash water run off to nearby ditch	Drainage ditches	Land	Wash water from poultry houses directed in sealed system to underground storage tank. Spillages of litter on yard areas during cleanout swept up, Lightly contaminated yard wash directed to underground tank. All clean site drainage directed to soakaways.	Unlikely	Pollution of watercourses leading to eutrophication and poisoning of flora and fauna	Not significant if managed carefully.
Pests						
Flies	Neighbouring dwelling houses within 400m of installation	Air	Temporary field heaps regularly checked for maggots and flies, heaps treated with pesticide and covered if flies become a an issue	Unlikely	Flies are a vector of pollution that can harm human health and amenity causing offence.	Not significant if managed carefully.

Rodents/Vermin	Neighbouring dwelling houses within 400m of installation	Land	Feed spillages cleared up promptly. Specialist contractor used to control pests.	Unlikely	Rodents are a vector of pollution that can harm human health and amenity causing offence.	Not significant if managed carefully.
----------------	--	------	--	----------	---	---------------------------------------